

REMARKS

The specification has been amended to provide a cross-reference to the previously filed International Application.

Claim 11 has been amended to remove the improper multi-dependency.

Entry of the Preliminary Amendment and favorable action on the merits are respectfully requested.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By *my/Well #36,623*
Andrew D. Meikle, #32,868

ADM/ja:cqc

P.O. Box 747
Falls Church, VA 22040-0747
(703) 205-8000

Attachment: Version With Markings Showing Changes Made

VERSION WITH MARKINGS SHOWING CHANGES MADE

The claims have been amended as follows:

11. (Amended) An antigen comprising the recombinant protein from merozoite of *Babesia caballi* as set forth in [any of claims 4 to 6] claim 4.

DESCRIPTION

GENE ENCODING PROTEIN FROM MEROZOITE OF *BABESIA CABALLI*,
RECOMBINANT PROTEIN OBTAINED WITH SAID GENE AND USE THEREOF
TECHNICAL FIELD

5 The present invention relates to a protein
derived from a merozoite of *Babesia caballi* (hereinafter
also referred to as "BC"), a kind of equine Protozoa
Babesia, a gene encoding said protein, an antibody specific
to said protein, and a method for diagnosing equine
10 babesiasis using the same.

BACKGROUND ART

Equine babesiasis is protoziasis carried by the
mites. The pathogen of this disease is equine Protozoa
Babesia, among which two species of *Babesia caballi* and
15 *Babesia equi* (hereinafter also referred to as "BE") are
known.

Equine babesiasis is widely spread all over the
world including South Europe, Asia, Russia, the Middle and
Near East, Africa, and Central and South America.
20 Clinically, this disease has main symptoms of anemia and
jaundice with high fever and progresses either acutely or
chronically. In acute cases, its lethality reaches about
10% or even as high as 50% in rare cases although it may
somewhat vary with either of the two pathogens. On the
25 other hand, the conditions after prognosis vary with either